

AD-A068 239

FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1  
FUNCTIONS DECREASING IN TRANSPOSITION WITH APPLICATIONS TO SHOC--ETC(U)  
MAR 79 M HOLLANDER, F PROSCHAN, J SETHURAMAN AFOSR-78-3678

UNCLASSIFIED

FSU-STATISTICS-M494

ARO-13888.18-MX

NL

| OF |

AD  
A068239




END

DATE

FILMED

6 --79

DDC



AD A068239

DDC FILE COPY

The Florida State University  
**Department  
of  
Statistics**  
Tallahassee, Florida



AD A068239

(14) FSU-STATISTICS-M494, TR-8

(12)

(9) Technical rept's

(6)

FUNCTIONS DECREASING IN TRANSPOSITION  
WITH APPLICATIONS TO SHOCK, DAMAGE, AND DOWN TIME.

by

Myles/Hollander\*, Frank/Proschan\* and Jayaram/Sethuraman†

(10)

FSU Statistics Report M494  
AFOSR Technical Report No. 8  
USARO-D No. 31

(13) 17 p.

(11)

March, 1979  
The Florida State University  
Department of Statistics  
Tallahassee, Florida

DDC FILE COPY

DDC  
RECEIVED  
MAY 2 1979  
C

(15)

✓ AFOSR-78-3678,  
✓ DAAG 29-76-G-0238

\* Research supported by the Air Force Office of Scientific Research, AFSC, USAF under Grant No. 78-3678.

† Research supported by the United States Army Research Office, Durham, under Grant No. DAA29-76-G-0238.

(18) ARD (19) 13888.18-MX

THE VIEW, OPINIONS, AND/OR FINDINGS CONTAINED IN THIS REPORT ARE THOSE OF THE AUTHOR(S) AND SHOULD NOT BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE ARMY POSITION, POLICY, OR DECISION, UNLESS SO DESIGNATED BY OTHER DOCUMENTATION.

This document has been approved for public release and sale; its distribution is unlimited.

400 277

Functions Decreasing in Transposition with Applications to  
Shock, Damage, and Down Time

M. Hollander\*, F. Proschan\*, and J. Sethuraman†

Florida State University, Tallahassee, Florida, USA

ACROSS	State Section	<input checked="" type="checkbox"/>
ALIS	Ref Section	<input type="checkbox"/>
BDC		
DATA/NOU/NO		
JUSTIFICATION		
BY	DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL	and/or SPECIAL
A		

In this paper a large class of multivariate densities and frequency functions, including the multivariate Poisson distribution and the compound multivariate Poisson distribution, are shown to have the decreasing in transposition property introduced by Hollander, Proschan, and Sethuraman (1977, Ann. Statist. 5, 722-733). Applications relevant to reliability are given, including applications to shock models, cumulation of damage, and component down times.

#### 1. INTRODUCTION AND PRELIMINARIES

In this paper we derive a basic theorem (Section 2, Theorem 2.1) showing that a large class of multivariate probability densities and frequency functions possess the decreasing in transposition (DT) property (Definition 1.3) introduced by Hollander, Proschan, and Sethuraman [3] (HPS[3]). Of particular interest are the multivariate Poisson distribution and the compound multivariate Poisson distribution treated in Section 3. Then, in Section 4, we consider applications arising in reliability. These applications pertain to shock models, cumulation of damage, and component down times. In the remainder of this section we give some preliminaries, including definitions and theorems, which will be useful in the sequel.

---

AMS 1970 subject classifications: Primary 62E99, 62H99; Secondary 62N05  
Key words and phrases: Functions decreasing in transposition, multivariate distributions, reliability, shock models.

\*Research supported by the Air Force Office of Scientific Research, AFSC, USAF under Grant No. 78-3678

†Research supported by the United States Army Research Office, Durham, under Grant No. DAA29-76-G-0238.

HPS[3] define a partial ordering on  $n$ -dimensional Euclidean space  $(R^n)$  presented in Definition 1.2 below.

DEFINITION 1.1. A vector  $\underline{x} = (x_1, \dots, x_n)$  is said to be a simple transposition of a vector  $\underline{x}'$  if  $\underline{x}$  and  $\underline{x}'$  agree in all but two coordinates, say  $i$  and  $j$ ,  $i < j$ ,  $x_i < x_j$ ,  $x'_i = x_j$ , and  $x'_j = x_i$ ; we write  $\underline{x} \overset{t}{\succ} \underline{x}'$ .

Thus  $\underline{x}'$  is obtained from  $\underline{x}$  by performing an inversion of a single pair of coordinates that occur in their natural order in  $\underline{x}$ .

DEFINITION 1.2. Let  $\underline{x}$  and  $\underline{x}'$  be two  $n$ -dimensional vectors such that there exists a finite number of vectors  $\underline{x}^0, \underline{x}^1, \dots, \underline{x}^k$  in  $R^n$  satisfying  $\underline{x} = \underline{x}^0 \overset{t}{\succ} \underline{x}^1 \overset{t}{\succ} \dots \overset{t}{\succ} \underline{x}^k = \underline{x}'$ ; i.e.,  $\underline{x}'$  is obtained from  $\underline{x}$  by a finite number of simple transpositions. We say that  $\underline{x}'$  is a transposition of  $\underline{x}$ .

HPS [3] then define a class of functions as follows:

DEFINITION 1.3. Let  $\underline{\lambda} = (\lambda_1, \dots, \lambda_n)$ , where  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$  are  $n$  ordered values in  $R^1$ . We say that  $f(\underline{\lambda}, \underline{x})$  is decreasing in transposition (DT) if:

(a)  $f(\underline{\lambda}^\pi, \underline{x}^\pi) = f(\underline{\lambda}, \underline{x})$  for each permutation  $\pi = (\pi_1, \dots, \pi_n)$  of the indices  $1, 2, \dots, n$ , where  $\underline{\lambda}^\pi = (\lambda_{\pi_1}, \dots, \lambda_{\pi_n})$  and

$$\underline{x}^\pi = (x_{\pi_1}, \dots, x_{\pi_n}).$$

(b)  $\underline{x} \overset{t}{\succ} \underline{x}'$  implies that  $f(\underline{\lambda}, \underline{x}) \geq f(\underline{\lambda}, \underline{x}')$ .

The familiar concepts of majorization and Schur functions will be used in this paper, and for completeness we recall their definitions.

DEFINITION 1.4. Let  $x_{[1]} \geq \dots \geq x_{[n]}$  be a decreasing rearrangement of the coordinates of the vector  $\underline{x}$ . Let  $\underline{x}$  and  $\underline{y}$  satisfy:

$$\sum_{i=1}^j x_{[i]} \geq \sum_{i=1}^j y_{[i]}, \quad j = 1, \dots, n-1$$

$$\sum_{i=1}^n x_{[i]} = \sum_{i=1}^n y_{[i]}.$$

Then  $\underline{x}$  is said to majorize  $\underline{y}$  (we write  $\underline{x} \stackrel{m}{\geq} \underline{y}$ ).

DEFINITION 1.5. A function  $f$  from  $R^n$  into  $R^1$  is said to be Schur-convex (Schur-concave) if  $\underline{x} \stackrel{m}{\geq} \underline{y}$  implies  $f(\underline{x}) \geq (\leq) f(\underline{y})$ .

HPS[3] derive basic properties of DT functions and show their relationship to other classes of functions. They show (HPS[3], Lemma 2.2) that when  $f(\underline{\lambda}, \underline{x})$  is of the form  $g(\underline{\lambda} - \underline{x})$ ,  $f(\underline{\lambda}, \underline{x})$  is a DT function if and only if  $g(\underline{y})$  is a Schur-concave function. They also show (HPS[3], Theorem 3.2) that the DT property is preserved under positive mixtures and (HPS[3], Theorem 3.6) that the DT property is preserved under products of positive DT functions. They also prove a "composition" theorem for DT functions (HPS[3], Theorem 3.3) and a "preservation" theorem (HPS[3], Theorem 3.7) for Schur-concave functions under an integral transform where the kernel is DT.

By restricting  $g_2(\underline{y}, \underline{z})$  in the HPS "composition" theorem to be of the form  $g(\underline{y} - \underline{z})$ , where  $g(\underline{w})$  is Schur-concave, we have the following.

THEOREM 1.5. Let  $g_1(\underline{x}, \underline{y})$  be a DT function and  $g(\underline{w})$  be a Schur-concave function such that

$$f(\underline{x}, \underline{z}) = \int g_1(\underline{x}, \underline{y}) g(\underline{y} - \underline{z}) d\mu(\underline{y})$$

is well-defined, where  $\mu$  is a positive permutation invariant measure. Then  $f(\underline{x}, \underline{z})$  is a DT function.

Theorem 1.5 generalizes the following result of Marshall and Olkin ([5], Theorem 2.1).

THEOREM 1.6. (Marshall-Olkin, [5]). Let  $g_1(\underline{x})$  and  $g_2(\underline{x})$  be Schur-concave functions such that

$$f(\underline{x}) = \int g_1(\underline{x} - \underline{y}) g_2(\underline{y}) d\mu(\underline{y})$$

is well-defined, where  $\mu$  is as in Theorem 1.5. Then  $f(\underline{x})$  is a Schur-concave function.

## 2. DT PROPERTY OF OVERLAPPING SUMS

In this section we state and prove our main theorem. Let, for  $k = 2, 3, \dots, n$ ,

$$\Lambda_k = \{\lambda: \lambda \text{ is a subset of size } k \text{ from } \{1, \dots, n\}\}. \quad (2.1)$$

THEOREM 2.1. Let  $\underline{X} = (X_1, \dots, X_n)$ ,  $\{X_\lambda, \lambda \in \Lambda_k\}$ ,  $k = 2, \dots, n$ , be independent collections of random variables. Let  $\underline{X}$  have a DT density function. Let the random variables in  $\{X_\lambda, \lambda \in \Lambda_k\}$  be i.i.d. and have a common log-concave density function  $g_k$ ,  $k = 2, \dots, n$ . Let, for  $i = 1, \dots, n$ ,

$$Z_i = X_i + \sum_{k=2}^i \sum_{\lambda: \lambda \in \Lambda_k \text{ and } i \in \lambda} X_\lambda. \quad (2.2)$$

Then  $\underline{Z} = (Z_1, \dots, Z_n)$  has a DT density function.

REMARK 2.2. Note that the summands appearing in the  $Z_1, \dots, Z_n$  overlap considerably. For example,  $X_{12}$  appears in the expressions for  $Z_1$  and  $Z_2$ ,  $X_{123}$  appears in the expressions for  $Z_1, Z_2$ , and  $Z_3$ , etc. Thus the inheritance of the DT property of  $\underline{Z}$  from that of  $\underline{X}$  is complicated by the overlapping of the  $X$ 's.

To prove the main result, we shall find it helpful to have available the following lemma:

LEMMA 2.3 Let  $k \geq 2$ . Let  $\{X_\lambda, \lambda \in \Lambda_k\}$  be i.i.d. random variables with a common log-concave density function  $g$  (with respect to the counting measure on a lattice or the Lebesgue measure). Let

$$W_i = \sum_{\lambda: \lambda \in \Lambda_k \text{ and } i \in \lambda} X_\lambda, \quad i = 1, \dots, n. \quad (2.3)$$

Let  $f(w_1, \dots, w_n)$  be the density function of  $\underline{W} = (w_1, \dots, w_n)$ . Then  $f$  is a Schur-concave function.

Proof. We will now give a proof for the case where  $g$  and  $f$  are density functions with respect to counting measures. The proof for the case where  $g$  and  $f$  are density functions with respect to Lebesgue measures is similar and will be omitted.

Notice first that  $W_1, \dots, W_n$  are exchangeable and hence  $f(w_1, \dots, w_n)$  is permutation invariant. Fix  $w_3, \dots, w_n$  and define

$$A_{w_1, w_2} = \{W_1 = w_1, W_2 = w_2, W_3 = w_3, \dots, W_n = w_n\}.$$

To show that  $f$  is Schur-concave, we must show that

$$P(A_{w_1, w_2}) \leq P(A_{w'_1, w'_2}) \quad (2.4)$$

whenever

$$(w_1, w_2) \succeq (w'_1, w'_2),$$

i.e. whenever

$$0 \leq \beta \leq \alpha \leq w^* \quad (2.5)$$

where

$$w_1 = w^* + \alpha, w_2 = w^* - \alpha, w'_1 = w^* + \beta, w'_2 = w^* - \beta.$$

The following classes of subsets of  $\{1, \dots, n\}$ , will help us to divide the elements of  $\Lambda$  into those that contain both 1 and 2, and those that do not contain either 1 or 2, and those that contain exactly one of 1 and 2. Define

$$V = \{v: v \text{ is a subset of size } (k-2) \text{ from } \{3, \dots, n\}\},$$

$$U = \{u: u \text{ is a subset of size } (k-1) \text{ from } \{3, \dots, n\}\},$$

and

$$S = \{s: s \text{ is a subset of size } k \text{ from } \{3, \dots, n\}\}.$$

Any  $\lambda$  in  $\Lambda_k$  is of the form  $12v$ ,  $1u$ ,  $2u$ , or  $s$ . Fix numbers  $x_{12v}$ ,  $v \in V$ ,  $x_s$ ,  $s \in S$  and  $x_u$ ,  $u \in U$  and define the event  $B$  as follows:

$$B = \{X_{12v} = x_{12v}, v \in V, X_s = x_s, s \in S, X_{1u} + X_{2u} = 2x_u, u \in U\}. \quad (2.6)$$

Let

$$2Y_u = X_{1u} - X_{2u}, u \in U.$$

Then the density function of  $\{Y_u, u \in U\}$  conditional on  $B$  is

$$\frac{\prod_u f(x_u + y_u) f(x_u - y_u)}{\sum_{y_u, u \in U} \prod_u [f(x_u + y_u) f(x_u - y_u)]}, \quad (2.7)$$

and the conditional probability of  $A_{w_1, w_2}$  given  $B$  is

$$P(A_{w_1, w_2} | B) = \begin{cases} \frac{\sum_{y_u, u \in U} \prod_u f(x_u + y_u) f(x_u - y_u)}{\sum_{y_u, u \in U} \prod_u [f(x_u + y_u) f(x_u - y_u)]} & \text{if } w_j = \sum_{v: j \in v} x_{12v} + \sum_{u: j \in u} x_u + \sum_{s: j \in s} x_s, j = 3, \dots, n \\ \text{and } w^* = \sum_v x_{12v} + \sum_u x_u & \\ 0 & \text{otherwise.} \end{cases}$$

Now, for each  $x_u$ ,  $f(x_u + y_u) f(x_u - y_u)$  is symmetric and log-concave in  $y_u$ .

The numerator in the expression for  $P(A_{w_1, w_2} | B)$  is therefore a convolution

of symmetric log-concave densities and hence, from Theorem 1 of Karlin and Proschan [4], is symmetric and log-concave in  $\alpha$ . Thus if (2.5) is satisfied, we obtain

$$P(A_{w_1, w_2} | B) \leq P(A_{w_1', w_2'} | B).$$

Inequality (2.4) now follows by unconditioning, and thus we have established that  $f$  is schur-concave. ||

We now give the proof of the main theorem.

Proof of Theorem 2.1. Let

$$X_i^{(k)} = \sum_{\lambda: \lambda \in \Lambda_k \text{ and } i \in \lambda} X_\lambda, \quad i = 1, \dots, n, k = 2, \dots, n.$$

Let  $\underline{X}^{(k)} = (X_1^{(k)}, \dots, X_n^{(k)})$ ,  $k = 2, \dots, n$ , and  $\underline{X} = (X_1, \dots, X_n)$ . Then

$$\underline{Z} = \underline{X} + \underline{X}^{(2)} + \dots + \underline{X}^{(n)}.$$

From Lemma 2.3, the density functions of  $\underline{X}^{(2)}, \dots, \underline{X}^{(n)}$  are all Schur-concave. Since the density function of  $\underline{Z}$  is the convolution of a DT density and several Schur-concave densities, it follows from Theorems 1.6 and 1.7 that the density function of  $\underline{Z}$  is DT. ||

### 3. APPLICATIONS TO MULTIVARIATE POISSON DISTRIBUTIONS

In this section we present several applications of Theorem 2.1.

3.1. MULTIVARIATE POISSON. The multivariate Poisson distribution (cf. Teicher [5], Dwass and Teicher [1]) can be defined as follows. Let  $X_i$  be a Poisson random variable with mean  $m_i$ ,  $i = 1, \dots, n$ ,  $X_\lambda$  be a Poisson random variable with mean  $m_\lambda$ ,  $\lambda \in \Lambda_k$ ,  $k = 2, \dots, n$ , where  $\Lambda_k$  is

given by (2.1). All the Poisson random variables are assumed mutually independent. Then  $(Z_1, \dots, Z_n)$ , as defined by (2.2), is said to have a joint multivariate Poisson distribution.

Now we specialize further to the case where

$$m_\lambda = m_k^*, \lambda \in \Lambda_k, k = 2, \dots, n. \quad (3.1)$$

That is, all  $X$ 's with the same number ( $> 1$ ) of subscripts, have a common mean. Then, from Theorem 2.1, we conclude that the corresponding frequency function is decreasing in transposition. That is, let  $\underline{m} = (m_1, \dots, m_n)$  and let  $f(\underline{m}, \underline{z})$  denote the joint probability mass function of  $Z_1, \dots, Z_n$  (where the other means are held fixed as in (3.1)). Then when  $m_1 \leq \dots \leq m_n$ , and  $\underline{z} \preceq \underline{z}'$ , we have  $f(\underline{m}, \underline{z}) \geq f(\underline{m}, \underline{z}')$ .

3.2. COMPOUND MULTIVARIATE POISSON. The compound multivariate Poisson distribution described below generalizes the compound bivariate Poisson distribution considered by Holgate [2] in the context of certain ecological situations. Let  $Z_1, Z_2, \dots, Z_n$  denote the number of individuals of type 1, 2, ..., n, respectively, in a quadrat of land. We suppose that these individuals arise from independent clusters and assume that the number of clusters  $N$  is a Poisson random variable with parameter  $\lambda$ . In cluster  $j$ , there are

$$Z_i^j = X_i^j + \sum_{k=2}^n \sum_{\lambda: \lambda \in \Lambda_k \text{ and } i \in \lambda} X_\lambda^i, \quad ,$$

individuals of type  $i$ ,  $i = 1, \dots, n$ ,

where  $X_1^j, X_\lambda^j, \lambda \in \Lambda_k, k = 2, \dots, n$ , are independent Poisson random variables with parameters  $m_i, m_\lambda, k = 2, \dots, n$ , respectively, and satisfying  $m_\lambda = m_k^*$  for  $\lambda \in \Lambda_k, k = 2, \dots, n$ , for each cluster  $j$ . We have already stated that the clusters are assumed to be independent. Thus  $\underline{Z}^j = (Z_1^j, \dots, Z_n^j)$  has a multivariate Poisson distribution. The vector  $\underline{Z}$  of the number of individuals of the different types in the quadrat is given by

$$\underline{Z} = \underline{Z}^1 + \dots + \underline{Z}^N$$

and is said to have a compound multivariate Poisson distribution. We now show that the density of  $\underline{Z}$  is DT. Conditional on  $N = N_0, \underline{Z}$  is the sum of  $N_0$  i.i.d. multivariate Poisson vectors and therefore is multivariate Poisson from the infinite divisibility of the multivariate Poisson established by Dwass and Teicher [1]. Thus the conditional density of  $\underline{Z}$ , given  $N = N_0$ , is DT. By unconditioning, the density of  $\underline{Z}$  is also DT. This proof also shows that the random number of clusters  $N$ , could be any random variable taking values on the positive integers.

#### 4. APPLICATIONS TO RELIABILITY PROBLEMS

In this section we show sample applications of Theorem 2.1 in some commonly occurring reliability models.

4.1. NUMBER OF SHOCKS EXPERIENCED BY COMPONENTS OF SYSTEM. Consider a system of  $n$  components experiencing successive shocks of various types over time, each shock affecting one or more components. More speci-

fically, in a fixed interval of time, let  $X_i$  be the number of shocks affecting component  $i$  alone ( $i = 1, 2, \dots, n$ ),  $X_{ij}$  the number of shocks affecting components  $i$  and  $j$  simultaneously and no other components ( $1 \leq i < j \leq n$ ), ...,  $X_{12\dots n}$  the number of shocks affecting all  $n$  components simultaneously. Let  $(Z_1, \dots, Z_n)$  be defined as in (2.2). Then  $Z_i$  denotes the total number of shocks experienced by component  $i$ ,  $i = 1, \dots, n$ . Assume that  $(X_1, \dots, X_n)$ ,  $\{X_\lambda, \lambda \in \Lambda_k\}$ ,  $k = 2, \dots, n$  satisfy the conditions of Theorem 2.1. Then we conclude that the joint frequency function of  $(Z_1, \dots, Z_n)$  is DT.

4.2. CUMULATION OF DAMAGE. We consider a model similar to that of 4.1. However, now we study the damage accumulated by each component as a result of the shocks simultaneously affecting the various subsets of components. Let  $D_i$  be the damage accumulated by component  $i$  as a result of shocks affecting component  $i$  alone ( $i = 1, \dots, n$ ),  $D_{ij}$  be the damage accumulated by components  $i$  and  $j$  as a result of shocks affecting components  $i$  and  $j$  simultaneously, but no other components ( $1 \leq i < j \leq n$ ), ..., and  $D_{12\dots n}$  be the damage accumulated by each of the components as a result of shocks affecting all components simultaneously. Assume that the joint distribution of  $\underline{D} = (D_1, \dots, D_n)$  has a DT density function, that  $D_\lambda, \lambda \in \Lambda_k$  has a common log concave density function  $f_k$ ,  $k = 2, \dots, n$ . Assume further that  $\underline{D}; \{D_\lambda, \lambda \in \Lambda_k\}$ ,  $k = 2, \dots, n$  are mutually dependent.


Define:

$$T_i = D_i + \sum_{k=2} \sum_{\lambda: \lambda \in \Lambda_k, i \in \lambda} D_\lambda, \quad i = 1, \dots, n.$$

$T_i$  represents the total damage to component  $i$  accruing from all shocks affecting component  $i$  (and possibly affecting other components simultaneously),  $i = 1, \dots, n$ .

From Theorem 2.1 we conclude that the joint probability density of  $T_1, \dots, T_n$  is DT.

4.3. COMPONENT DOWN TIMES. Suppose that an  $n$ -component system is subject to repair (replacement). If any component fails, the system fails and the repair (replacement) time for that component is recorded. Likewise, if any subset of components fails simultaneously, the system fails, and the time it takes to repair (replace) each of the components in that subset is recorded. For example, the operating record of a two-component system is exhibited in Fig. 4.1 below.



4.1. Operating Record of a Two-Component System

Figure 4.1 indicates that a total of 4 repair (replacement) periods occurred during  $[0, t]$ : 1 period was due to the failure of component 1 alone, 1 period was due to the failure of component 2 alone, and 2 periods were due to the simultaneous failure of components 1 and 2.

Let  $R_i$  = time devoted to repair of component  $i$  due to shocks affecting component  $i$  alone ( $1 \leq i \leq n$ ),  $R_\lambda$  = time devoted to repair of each of components in  $\lambda$  due to shocks affecting components in  $\lambda$  simultaneously and no other components, for each  $\lambda \in \Lambda_k$ ,  $k = 2, \dots, n$ . Assume that the joint distribution of  $R_1, \dots, R_n$  has a DT density function, that  $R_\lambda$  has a common log concave density function  $f_k$  for  $\lambda \in \Lambda_k$ ,  $k = 2, \dots, n$ . Assume further that  $\underline{R}, \{R_\lambda, \lambda \in \Lambda_k\}, k = 2, \dots, n$  are mutually independent. Define

$$S_i = R_i + \sum_{k=2}^n \sum_{\lambda: \lambda \in \Lambda_k, i \in \lambda} R_\lambda, \quad i = 1, \dots, n.$$

$S_i$  represents the total time spent repairing (replacing) component  $i$  as a result of shocks affecting component  $i$ ,  $i = 1, \dots, n$ .

From Theorem 2.1, we conclude that the joint density function of  $S_1, \dots, S_n$  is DT.

REFERENCES

- [1] Dwass, M. and Teicher, H. (1957). On infinitely divisible random vectors. Ann. Math. Statist. 28, 461-470.
- [2] Holgate, P. (1966). Bivariate generalizations of Neyman's Type A distribution. Biometrika 53, 241-245.
- [3] Hollander, M., Proschan, F., and Sethuraman, J. (1977). Functions decreasing in transposition and their applications in ranking problems. Ann. Statist. 5, 722-733.
- [4] Karlin, S. and Proschan, F. (1960). Pólya type distributions of convolutions. Ann. Math. Statist. 31, 721-726.
- [5] Marshall, A. W. and Olkin, I. (1974). Majorization in multivariate distributions. Ann. Statist. 2, 1189-1200.
- [6] Teicher, H. (1954). On the multivariate Poisson distribution. Skandinavisk Aktuarietidskrift 37, 1-9.

## UNCLASSIFIED

## Security Classification of this Page

## REPORT DOCUMENTATION PAGE

1. REPORT NUMBERS AFOSR 78-3678 No. 8 USARO-D-DAA29-76-G-0238 No. 31	2. GOVT. ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE FUNCTIONS DECREASING IN TRANSPOSITION WITH APPLICATIONS TO RELIABILITY	5. TYPE OF REPORT Technical Report	6. PERFORMING ORGANIZATION REPORT FSU Statistical Report No. M494
7. AUTHOR(s) Myles Hollander, Frank Proschan, and Jayaram Sethuraman	8. CONTRACT OR GRANT NUMBER(s) AFOSR 76-3678 No. 8 USARO-D-DAA29-76-G-0238 No. 31	
9. PERFORMING ORGANIZATION NAME & ADDRESS The Florida State University Department of Statistics Tallahassee, Florida 32306	10. PROGRAM ELEMENT, PROJECT, TASK AREA AND WORK UNIT NOS.	
11. CONTROLLING OFFICE NAME & ADDRESS Air Force Office of Scientific Research Bolling Air Force Base, D. C. 20332 United States Army Research Office, Durham, North Carolina	12. REPORT DATE February, 1979	13. NUMBER OF PAGES 14
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS UNCLASSIFIED	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release: distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS Functions decreasing in transposition, multivariate distributions, reliability, shock models		
20. ABSTRACT In this paper a large class of multivariate densities and frequency functions including the multivariate Poisson distribution and the compound multivariate Poisson distribution, are shown to have the decreasing in transposition property introduced by Hollander, Proschan, and Sethuraman (1977, <u>Ann. Statist.</u> 5, 722-733). Applications relevant to reliability are given, including applications to shock models, cumulation of damage, and component down times.		